



PB-210-T025

Monoclonal Antibody to CD11a Pacific Blue™ conjugated (25 tests)

Clone:	MEM-25
Isotype:	Mouse IgG1
Specificity:	The antibody MEM-25 reacts with CD11a (alpha subunit of human LFA-1), a 170-180 kDa type I transmembrane glycoprotein expressed on B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. HLDA IV; WS Code NL 209
Immunogen:	Leukocytes from a patient suffering from a LGL-type leukaemia.
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Pacific Blue [®] 488; under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity conformation. Cell stimulation by chemokines or other signals leads to induction the high-affinity conformation, which supports tight binding of LFA-1 to its ligands, the intercellular adhesion molecules ICAM-1, -2, -3. LFA-1 is thus involved in interaction of various immune cells and in their tissue-specific settlement, but participates also in control of cell differentiation and proliferation and of T-cell effector functions. Blocking of LFA-1 function by specific antibodies or small molecules has become an important therapeutic approach in treatment of multiple inflammatory diseases. For example, humanized anti-LFA-1 antibody Efalizumab (Raptiva) is being used to interfere with T cell migration to sites of inflammation; binding of cholesterol-lowering drug simvastatin to CD11a allosteric site leads to immunomodulation and increase in lymphocytic cholinergic activity.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Sarantos MR, Raychaudhuri S, Lum AF, Staunton DE, Simon SI: Leukocyte function-associated antigen 1-mediated adhesion stability is dynamically regulated through affinity and valency during bond formation with intercellular adhesion molecule-1. *J Biol Chem.* 2005 Aug 5;280(31):28290-8.
- *Giblin PA, Lemieux RM: LFA-1 as a key regulator of immune function: approaches toward the development of LFA-1-based therapeutics. *Curr Pharm Des.* 2006;12(22):2771-95.
- *Kellersch B, Kolanus W: Membrane-proximal signaling events in beta-2 integrin activation. *Results Probl Cell Differ.* 2006;43:245-57.
- *Fujii T, Takada-Takatori Y, Kawashima K: Roles played by lymphocyte function-associated antigen-1 in the regulation of lymphocytic cholinergic activity. *Life Sci.* 2007 May 30;80(24-25):2320-4.
- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- *Bazil V, Stefanova I, Hilgert I, Kristofova H, Vanek S, Horejsi V.: Monoclonal antibodies against human leucocyte antigens. IV. Antibodies against subunits of the LFA-1 (CD11a/CD18) leucocyte-adhesion glycoprotein. *Folia Biol (Praha).* 1990;36(1):41-50.
- *Otonello L, Epstein AL, Mancini M, Dapino P, Dallegri F: Monoclonal LYM-1 antibody-dependent cytotoxicity by human neutrophils exposed to GM-CSF: auto-regulation of target cell attack by cathepsin G. *J Leukoc Biol.* 2004 Jan;75(1):99-105.
- *Simon SI, Cherapanov V, Nadra I, Waddell TK, Seo SM, Wang Q, Doerschuk CM, Downey GP: Signaling functions of L-selectin in neutrophils: alterations in the cytoskeleton and colocalization with CD18. *J Immunol.* 1999 Sep 1;163(5):2891-901.
- *Hajishengallis G, Martin M, Sojar HT, Sharma A, Schifferle RE, DeNardin E, Russell MW, Genco RJ: Dependence of bacterial protein adhesins on toll-like receptors for proinflammatory cytokine induction. *Clin Diagn Lab Immunol.* 2002 Mar;9(2):403-11.
- *Wang JH, Kwas C, Wu L: Intercellular adhesion molecule 1 (ICAM-1), but not ICAM-2 and -3, is important for dendritic cell-mediated human immunodeficiency virus type 1 transmission. *J Virol.* 2009 May;83(9):4195-204.
- *Garnotel R, Rittié L, Poitevin S, Monboisse JC, Nguyen P, Potron G, Maquart FX, Randoux A, Gillery P: Human blood monocytes interact with type I collagen through alpha x beta 2 integrin (CD11c-CD18, gp150-95). *J Immunol.* 2000 Jun 1;164(11):5928-34.
- *Mathison RD, Befus AD, Davison JS, Woodman RC: Modulation of neutrophil function by the tripeptide feG. *BMC Immunol.* 2003 Mar 4;4:3. Epub 2003 Mar 4.
- *Otonello L, Epstein AL, Dapino P, Barbera P, Morone P, Dallegri F: Monoclonal Lym-1 antibody-dependent cytotoxicity by neutrophils exposed to granulocyte-macrophage colony-stimulating factor: intervention of Fc gamma RII (CD32), CD11b-CD18 integrins, and CD66b glycoproteins. *Blood.* 1999 May 15;93(10):3505-11.
- *Aubert M, Yoon M, Sloan DD, Spear PG, Jerome KR: The virological synapse facilitates herpes simplex virus entry into T cells. *J Virol.* 2009 Jun;83(12):6171-83.
- *And many other.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz