



11-213-C100

Monoclonal Antibody to CD15 Purified Antibody (0.1 mg)

Clone:	MEM-158
Isotype:	Mouse IgM
Specificity:	The antibody MEM-158 reacts with CD15, a cell membrane molecule 3-fucosyl-N-acetyllactosamine (3-FAL) strongly expressed on granulocytes, monocytes, macrophages, mast cells; it is also present on Langerhans cells and some myeloid precursors cells. HLDA VI; WS Code AS A053
Immunogen:	Human granulocytes
Species Reactivity:	Human
Negative Species:	Porcine
Application:	Flow Cytometry Immunoprecipitation
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified from ascites by precipitation methods and ion exchange chromatography.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not use after expiration date stamped on vial label. Do not freeze.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD15 (Lewis X, Le(x); stage specific embryonic antigen-1, SSEA-1) is a trisaccharide determinant (3-fucosyl-N-acetyllactosamine) expressed on several glycolipids, glycoproteins and proteoglycans of various cell types, e.g. granulocytes, mast cells, monocytes, macrophages, cells of gastric mucosa, nervous system or various tumour cells. There are several variants of Lewis x, such as sialyl-Lewis x or sulphated Lewis x. Cells with high surface expression of Le(x) antigen exhibit strong self-aggregation, based on calcium-dependent Le(x)-Le(x) interaction. This process is involved for example in embryo compaction or in autoaggregation of teratocarcinoma cells. Sialyl-Le(x) and its isomer sialyl-Le(a) are ligands of selectins. CD15 expression has been extensively used to confirm diagnosis of Hodgkin's disease.
References:	*Benharroch D, Dima E, Levy A, Ohana-Malka O, Ariad S, Prinsloo I, Mejirovsky E, Sacks M, Gopas J: Differential expression of sialyl and non-sialyl-CD15 antigens on Hodgkin-Reed-Sternberg cells: significance in Hodgkin's disease. <i>Leuk Lymphoma</i> . 2000 Sep;39(1-2):185-94. *Hakomori S: Le(X) and related structures as adhesion molecules. <i>Histochem J</i> . 1992 Nov;24(11):771-6. *Li C, Wong P, Pan T, Xiao F, Yin S, Chang B, Kang SC, Ironside J, Sy MS: Normal cellular prion protein is a ligand of selectins: binding requires Le(X) but is inhibited by sLe(X). <i>Biochem J</i> . 2007 Sep 1;406(2):333-41. *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

For laboratory research only, not for drug, diagnostic or other use.