

A7-214-T025

Monoclonal Antibody to CD16 Alexa Fluor® 700 conjugated (25 tests)

Clone:	MEM-154
Isotype:	Mouse IgG1
Specificity:	<p>The antibody MEM-154 reacts with the epitope on CD16 antigen that residing in proximity to FG loop (probably BC or C'E loop). CD16 is a low affinity receptor for aggregated IgG (FcγRIII antigen). The antibody MEM-154 reacts with CD16+ granulocytes.</p> <p>HLDA V; WS Code M MA068 HLDA V; WS Code NK NK51</p>
Immunogen:	Human granulocytes
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Alexa Fluor® 700 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	<p>Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.</p> <p>Do not use after expiration date stamped on vial label.</p> <p>Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.</p>
Usage:	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.</p> <p>The content of a vial (0.1 ml) is sufficient for 25 tests.</p>
Expiration:	See vial label
Lot Number:	See vial label
Background:	<p>CD16 (FcγRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcγRIII is expressed in two forms &#8211; FcγRIII-A and -B. FcγRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcεRI-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcγRIII-A is associated, moreover, with FcεRI-beta subunit. Besides IgG, FcγRIII-A can be triggered also by oligomeric IgE. FcγRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.</p>

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Gessner JE, Grussenmeyer T, Kolanus W, Schmidt RE: The human low affinity immunoglobulin G Fc receptor III-A and III-B genes. Molecular characterization of the promoter regions. *J Biol Chem.* 1995 Jan 20;270(3):1350-61.
- *Kocher M, Siegel ME, Edberg JC, Kimberly RP: Cross-linking of Fc gamma receptor IIa and Fc gamma receptor IIIb induces different proadhesive phenotypes on human neutrophils. *J Immunol.* 1997 Oct 15;159(8):3940-8.
- *Arase N, Arase H, Hirano S, Yokosuka T, Sakurai D, Saito T: IgE-mediated activation of NK cells through Fc gamma RIII. *J Immunol.* 2003 Mar 15;170(6):3054-8.
- *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- *de Haas M, Koene HR, Kleijer M, de Vries E, Simsek S, van Tol MJ, Roos D, von dem Borne AE: A triallelic Fc gamma receptor type IIIA polymorphism influences the binding of human IgG by NK cell Fc gamma RIIIa. *J Immunol.* 1996 Apr 15;156(8):3948-55.
- *Tamm A, Schmidt RE: The binding epitopes of human CD16 (Fc gamma RIII) monoclonal antibodies. Implications for ligand binding. *J Immunol.* 1996 Aug 15;157(4):1576-81.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.