

1P-221-T025

Monoclonal Antibody to CD44 Phycoerythrin (PE) conjugated (25 tests)

Clone:	MEM-85
Isotype:	Mouse IgG2b
Specificity:	<p>The antibody MEM-85 reacts with both cell surface-expressed and soluble form of CD44 antigen (Phagocyte glycoprotein 1), a 80-95 kDa transmembrane glycoprotein (hyaladherin family) present on the most of cells and tissues (leukocytes, endothelial cells, mesenchymal cells, etc.); it is negative on platelets and hepatocytes.</p> <p>HLDA IV; WS Code NL 706 HLDA VI; WS Code AS Ref.15</p>
Immunogen:	Leukocytes of a patient suffering from LGL Type Leukaemia.
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	<p>Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.</p> <p>Do not use after expiration date stamped on vial label.</p> <p>Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.</p>
Usage:	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.</p> <p>The content of a vial (0.5 ml) is sufficient for 25 tests.</p>
Expiration:	See vial label
Lot Number:	See vial label
Background:	<p>CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculation and homing, it can thus serve e.g. as a modulator of macrophage recruitment in response to pathogen. Although CD44 functions are essential for physiological activities of normal cells, elevated CD44 expression correlates with poor prognosis in many carcinomas, facilitating tumour growth and metastasis, antiapoptosis and directional motility of cancer cells.</p>

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Vigetti D, Viola M, Karousou E, Rizzi M, Moretto P, Genasetti A, Clerici M, Hascall VC, De Luca G, Passi A: Hyaluronan-CD44-ERK1/2 regulate human aortic smooth muscle cell motility during aging. *J Biol Chem.* 2007 Dec 12
- *Hollingsworth JW, Li Z, Brass DM, Garantzotis S, Timberlake SH, Kim A, Hossain I, Savani RC, Schwartz DA. CD44 regulates macrophage recruitment to the lung in lipopolysaccharide-induced airway disease. *Am J Respir Cell Mol Biol.* 2007 Aug;37(2):248-53.
- *Liu J, Bi G, Wen P, Yang W, Ren X, Tang T, Xie C, Dong W, Jiang G. Down-regulation of CD44 contributes to the differentiation of HL-60 cells induced by ATRA or HMBA. *Cell Mol Immunol.* 2007 Feb;4(1):59-63.
- *Subramaniam V, Gardner H, Jothy S: Soluble CD44 secretion contributes to the acquisition of aggressive tumor phenotype in human colon cancer cells. *Exp Mol Pathol.* 2007 Dec;83(3):341-6.
- *Subramaniam V, Vincent IR, Gardner H, Chan E, Dhamko H, Jothy S: CD44 regulates cell migration in human colon cancer cells via Lyn kinase and AKT phosphorylation. *Exp Mol Pathol.* 2007 Oct;83(2):207-15.
- *Stefanová I, Hilgert I, Bazil V, Kristofová H, Horejsí V: Human leucocyte surface glycoprotein CDw44 and lymphocyte homing receptor are identical molecules. *Immunogenetics.* 1989;29(6):402-4.
- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- *Bazil V, Strominger JL: Metalloprotease and serine protease are involved in cleavage of CD43, CD44, and CD16 from stimulated human granulocytes. Induction of cleavage of L-selectin via CD16. *J Immunol.* 1994 Feb 1;152(3):1314-22.
- *Ilangumaran S, Briol A, Hoessli DC: CD44 selectively associates with active Src family protein tyrosine kinases Lck and Fyn in glycosphingolipid-rich plasma membrane domains of human peripheral blood lymphocytes. *Blood.* 1998 May 15;91(10):3901-8.
- *Kolar GR, Mehta D, Pelayo R, Capra JD: A novel human B cell subpopulation representing the initial germinal center population to express AID. *Blood.* 2007 Mar 15;109(6):2545-52.
- *Schmidt D, Achermann J, Odermatt B, Breyman C, Mol A, Genoni M, Zund G, Hoerstrup SP: Prenatally fabricated autologous human living heart valves based on amniotic fluid derived progenitor cells as single cell source. *Circulation.* 2007 Sep 11;116(11 Suppl):I64-70.

For laboratory research only, not for drug, diagnostic or other use.