

A6-235-T100

Monoclonal Antibody to CD71 Alexa Fluor® 647 conjugated (100 tests)

Clone:	MEM-75
Isotype:	Mouse IgG1
Specificity:	<p>The antibody MEM-75 reacts with CD71 antigen (transferrin receptor), a 95 kDa type II homodimeric transmembrane glycoprotein expressed on activated B and T lymphocytes, macrophages and erythroid precursors; it is lost on resting blood leukocytes.</p> <p>The antibody MEM-75 does not block binding of transferrin to the receptor. HLDA IV; WS Code A 45 HLDA V; WS Code T T-165</p>
Immunogen:	NALM-6 human pre-B cell line
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Alexa Fluor® 647 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	<p>Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.</p> <p>Do not use after expiration date stamped on vial label.</p> <p>Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.</p>
Usage:	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.</p> <p>The content of a vial (0.4 ml) is sufficient for 100 tests.</p>
Expiration:	See vial label
Lot Number:	See vial label
Background:	<p>CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clathrin-mediated endocytosis. Acidification of endosomes by vesicular membrane proton pumps leads to dissociation of iron ions, whereas transferrin (apotransferrin) remains associated with CD71 and recycles to the cell surface, where it is released upon exposure to normal pH. CD71 is also involved in uptake of non-transferrin bound iron.</p>

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Rouault TA: How mammals acquire and distribute iron needed for oxygen-based metabolism. *PLoS Biol.* 2003 Dec;1(3):E79.
- *Taketani S: Aquisition, mobilization and utilization of cellular iron and heme: endless findings and growing evidence of tight regulation. *Tohoku J Exp Med.* 2005 Apr;205(4):297-318.
- *Graham RM, Chua AC, Herbison CE, Olynyk JK, Trinder D: Liver iron transport. *World J Gastroenterol.* 2007 Sep 21;13(35):4725-36.
- *Graham RM, Reutens GM, Herbison CE, Delima RD, Chua AC, Olynyk JK, Trinder D: Transferrin receptor 2 mediates uptake of transferrin-bound and non-transferrin-bound iron. *J Hepatol.* 2008 Feb;48(2):327-34.
- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- *Beck Z, Balogh A, Kis A, Izsépi E, Cervenak L, László G, Bíró A, Liliom K, Mocsár G, Vámosi G, Füst G, Matko J: New cholesterol-specific antibodies remodel HIV-1 target cells' surface and inhibit their in vitro virus production. *J Lipid Res.* 2010 Feb;51(2):286-96.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz