



Antibodies

1F-235-T100

## Monoclonal Antibody to CD71 Fluorescein (FITC) conjugated (100 tests)

<b>Clone:</b>	MEM-75
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	<p>The antibody MEM-75 reacts with CD71 antigen (transferrin receptor), a 95 kDa type II homodimeric transmembrane glycoprotein expressed on activated B and T lymphocytes, macrophages and erythroid precursors; it is lost on resting blood leukocytes.</p> <p>The antibody MEM-75 does not block binding of transferrin to the receptor. HLDA IV; WS Code A 45 HLDA V; WS Code T T-165</p>
<b>Immunogen:</b>	NALM-6 human pre-B cell line
<b>Species Reactivity:</b>	Human
<b>Preparation:</b>	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
<b>Storage / Stability:</b>	<p>Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.</p> <p>Do not use after expiration date stamped on vial label.</p> <p>Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.</p>
<b>Usage:</b>	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension.</p> <p>The content of a vial (2 ml) is sufficient for 100 tests.</p>
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	<p>CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clathrin-mediated endocytosis. Acidification of endosomes by vesicular membrane proton pumps leads to dissociation of iron ions, whereas transferrin (apotransferrin) remains associated with CD71 and recycles to the cell surface, where it is released upon exposure to normal pH. CD71 is also involved in uptake of non-transferrin bound iron.</p>

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

**References:**

- \*Rouault TA: How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 2003 Dec;1(3):E79.
- \*Taketani S: Aquisition, mobilization and utilization of cellular iron and heme: endless findings and growing evidence of tight regulation. Tohoku J Exp Med. 2005 Apr;205(4):297-318.
- \*Graham RM, Chua AC, Herbison CE, Olynyk JK, Trinder D: Liver iron transport. World J Gastroenterol. 2007 Sep 21;13(35):4725-36.
- \*Graham RM, Reutens GM, Herbison CE, Delima RD, Chua AC, Olynyk JK, Trinder D: Transferrin receptor 2 mediates uptake of transferrin-bound and non-transferrin-bound iron. J Hepatol. 2008 Feb;48(2):327-34.
- \*Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- \*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- \*Beck Z, Balogh A, Kis A, Izsépi E, Cervenak L, László G, Bíró A, Liliom K, Mocsár G, Vámosi G, Füst G, Matko J: New cholesterol-specific antibodies remodel HIV-1 target cells' surface and inhibit their in vitro virus production. J Lipid Res. 2010 Feb;51(2):286-96.

**For laboratory research only, not for drug, diagnostic or other use.**