



A4-292-C100

## Monoclonal Antibody to HLA-G Alexa Fluor® 488 conjugated (0.1 mg)

<b>Clone:</b>	MEM-G/9
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	<p>The antibody MEM-G/9 reacts with native form of human HLA-G1 on the cell surface as well as with soluble HLA-G5 isoform in its beta2-microglobulin associated form. HLA-G belongs to the MHC Class I molecules (MHC Class Ib; nonclassical) and it is expressed on the surface of trophoblast cells.</p> <p>The antibody MEM-G/9 is standard reagent thoroughly validated during 3rd International Conference on HLA-G (Paris, 2003).</p>
<b>Immunogen:</b>	Recombinant human HLA-G refolded with beta2-microglobulin and peptide.
<b>Species Reactivity:</b>	Human
<b>Negative Species:</b>	Mouse
<b>Preparation:</b>	The purified antibody is conjugated with Alexa Fluor 488 under optimum conditions. The conjugate is purified by size-exclusion chromatography.
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	<p>Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.</p> <p>Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.</p>
<b>Usage:</b>	<p>The reagent is designed for Flow Cytometry analysis.</p> <p>Suggested working dilution is 5 µg/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.</p>
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

- \*Fournel S, Huc X, Aguerre-Girr M, Solier C, Legros M, Praud-Brethenou C, Moussa M, Chaouat G, Berrebi A, Bensussan A, Lenfant F, Le Bouteiller P.: Comparative reactivity of different HLA-G monoclonal antibodies to soluble HLA-G molecules. *Tissue Antigens*. 2000 Jun;55(6):510-8.
- \*Lozano JM, Gonzalez R, Kindelan JM, Rouas-Freiss N, Caballos R, Dausset J, Carosella ED, Pena J.: Monocytes and T lymphocytes in HIV-1-positive patients express HLA-G molecule. *AIDS*. 2002 Feb 15;16(3):347-51.
- \*Pangault C, Le Fric G, Caulet-Maugendre S, Lena H, Amiot L, Guilloux V, Onno M, Fauchet R.: Lung macrophages and dendritic cells express HLA-G molecules in pulmonary diseases. *Hum Immunol*. 2002 Feb;63(2):83-90.
- \*Fuzzi B, Rizzo R, Criscuoli L, Noci I, Melchiorri L, Scarselli B, Bencini E, Menicucci A, Baricordi OR.: HLA-G expression in early embryos is a fundamental prerequisite for the obtainment of pregnancy. *Eur J Immunol*. 2002 Feb;32(2):311-5.
- \*Menier C, Saez B, Horejsi V, Martinuzzi S, Krawice-Radanne I, Bruel S, Le Danff C, Reboul M, Hilgert I, Rabreau M, Larrad ML, Pla M, Carosella ED, Rouas-Freiss N.: Characterization of monoclonal antibodies recognizing HLA-G or HLA-E: new tools to analyze the expression of nonclassical HLA class I molecules. *Hum Immunol*. 2003 Mar;64(3):315-26.
- \*Abstracts from the 3rd International Conference on HLA-G. *Tissue Antigens* 62, 339-357 (2003).
- \*Lopez AS, Alegre E, LeMaout J, Carosella E, Gonzalez A. Regulatory role of tryptophan degradation pathway in HLA-G expression by human monocyte-derived dendritic cells. *Mol Immunol*. 2006 Jul;43(14):2151-60.
- \*Gonen-Gross T, Achdout H, Arnon TI, Gazit R, Stern N, Horejsi V, Goldman-Wohl D, Yagel S, Mandelboim O: The CD85J/leukocyte inhibitory receptor-1 distinguishes between conformed and beta 2-microglobulin-free HLA-G molecules. *J Immunol*. 2005 Oct 15;175(8):4866-74.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz