



1P-292-C100

Monoclonal Antibody to HLA-G Phycoerythrin (PE) conjugated (0.1 mg)

Clone:	MEM-G/9
Isotype:	Mouse IgG1
Specificity:	<p>The antibody MEM-G/9 reacts with native form of human HLA-G1 on the cell surface as well as with soluble HLA-G5 isoform in its beta2-microglobulin associated form. HLA-G belongs to the MHC Class I molecules (MHC Class Ib; nonclassical) and it is expressed on the surface of trophoblast cells.</p> <p>The antibody MEM-G/9 is standard reagent thoroughly validated during 3rd International Conference on HLA-G (Paris, 2003).</p>
Immunogen:	Recombinant human HLA-G refolded with beta2-microglobulin and peptide.
Species Reactivity:	Human
Negative Species:	Mouse
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.1 mg/ml
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	<p>Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.</p> <p>Do not use after expiration date stamped on vial label.</p> <p>Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.</p>
Usage:	<p>The reagent is designed for Flow Cytometry analysis of cells expressing HLA-G molecule on the cell surface.</p> <p>Suggested working dilution is 1:100. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.</p>
Expiration:	See vial label
Lot Number:	See vial label

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Fournel S, Huc X, Aguerre-Girr M, Solier C, Legros M, Praud-Brethenou C, Moussa M, Chaouat G, Berrebi A, Bensussan A, Lenfant F, Le Bouteiller P.: Comparative reactivity of different HLA-G monoclonal antibodies to soluble HLA-G molecules. *Tissue Antigens*. 2000 Jun;55(6):510-8.
- *Lozano JM, Gonzalez R, Kindelan JM, Rouas-Freiss N, Caballos R, Dausset J, Carosella ED, Pena J.: Monocytes and T lymphocytes in HIV-1-positive patients express HLA-G molecule. *AIDS*. 2002 Feb 15;16(3):347-51.
- *Pangault C, Le Fric G, Caulet-Maugendre S, Lena H, Amiot L, Guilloux V, Onno M, Fauchet R.: Lung macrophages and dendritic cells express HLA-G molecules in pulmonary diseases. *Hum Immunol*. 2002 Feb;63(2):83-90.
- *Fuzzi B, Rizzo R, Criscuoli L, Noci I, Melchiorri L, Scarselli B, Bencini E, Menicucci A, Baricordi OR.: HLA-G expression in early embryos is a fundamental prerequisite for the obtainment of pregnancy. *Eur J Immunol*. 2002 Feb;32(2):311-5.
- *Menier C, Saez B, Horejsi V, Martinozzi S, Krawice-Radanne I, Bruel S, Le Danff C, Reboul M, Hilgert I, Rabreau M, Larrad ML, Pla M, Carosella ED, Rouas-Freiss N.: Characterization of monoclonal antibodies recognizing HLA-G or HLA-E: new tools to analyze the expression of nonclassical HLA class I molecules. *Hum Immunol*. 2003 Mar;64(3):315-26.
- *Abstracts from the 3rd International Conference on HLA-G. *Tissue Antigens* 62, 339-357 (2003).
- *Lopez AS, Alegre E, LeMaout J, Carosella E, Gonzalez A. Regulatory role of tryptophan degradation pathway in HLA-G expression by human monocyte-derived dendritic cells. *Mol Immunol*. 2006 Jul;43(14):2151-60.
- *Gonen-Gross T, Achdout H, Arnon TI, Gazit R, Stern N, Horejsi V, Goldman-Wohl D, Yagel S, Mandelboim O: The CD85J/leukocyte inhibitory receptor-1 distinguishes between conformed and beta 2-microglobulin-free HLA-G molecules. *J Immunol*. 2005 Oct 15;175(8):4866-74.

For laboratory research only, not for drug, diagnostic or other use.