



Antibodies

A6-237-C100

Monoclonal Antibody to beta2-microglobulin Alexa Fluor® 647 conjugated (0.1 mg)

Clone:	B2M-01
Isotype:	Mouse IgG2a
Specificity:	The antibody B2M-01 reacts with beta2-microglobulin (beta2M) associated with cell-surface MHC Class I molecules and other membrane antigens as well as with soluble beta2-microglobulin. Beta2M is a 12 kDa Ig like glycoprotein expressed on lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells and epithelial cells. It is absent on erythrocytes.
Immunogen:	Purified human beta2-microglobulin
Species Reactivity:	Human
Negative Species:	Mouse, Bovine, Canine (Dog), Rabbit, Chicken
Preparation:	The purified antibody is conjugated with Alexa Fluor 647 under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.
Usage:	The reagent is designed for Flow Cytometry analysis. Suggested working dilution is 2 µg/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Beta2-microglobulin non-covalently associates with the 44 kDa alpha chain to forms the HLA Class I antigen complex. Human beta2-microglobulin associated with HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes.
References:	*Hilgert I, Horejsi V, Kristofova H.: The use of murine monoclonal antibody B2M-01 for detection and purification of human beta 2-microglobulin. Folia Biol (Praha). 1984;30(6):369-76. *Khurana M, Traum AZ, Aivado M, Wells MP, Guerrero M, Grall F, Libermann TA, Schachter AD.: Urine proteomic profiling of pediatric nephrotic syndrome. Pediatr Nephrol. 2006 Sep;21(9):1257-65. Epub 2006 Jun 30.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 366 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz