



Antibodies

PB-237-C025

## Monoclonal Antibody to beta2-microglobulin Pacific Blue™ conjugated (0.025 mg)

<b>Clone:</b>	B2M-01
<b>Isotype:</b>	Mouse IgG2a
<b>Specificity:</b>	The antibody B2M-01 reacts with beta2-microglobulin (beta2M) associated with cell-surface MHC Class I molecules and other membrane antigens as well as with soluble beta2-microglobulin. Beta2M is a 12 kDa Ig like glycoprotein expressed on lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells and epithelial cells. It is absent on erythrocytes.
<b>Immunogen:</b>	Purified human beta2-microglobulin
<b>Species Reactivity:</b>	Human
<b>Negative Species:</b>	Mouse, Bovine, Canine (Dog), Rabbit, Chicken
<b>Preparation:</b>	The purified antibody is conjugated with Pacific Blue under optimum conditions. The conjugate is purified by size-exclusion chromatography.
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not use after expiration date stamped on vial label. Do not freeze.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis. Suggested working dilution is 1:300. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	<b>Beta2-microglobulin</b> non-covalently associates with the 44 kDa alpha chain to forms the HLA Class I antigen complex. Human beta2-microglobulin associated with HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes.
<b>References:</b>	*Hilgert I, Horejsi V, Kristofova H.: The use of murine monoclonal antibody B2M-01 for detection and purification of human beta 2-microglobulin. Folia Biol (Praha). 1984;30(6):369-76. *Khurana M, Traum AZ, Aivado M, Wells MP, Guerrero M, Grall F, Libermann TA, Schachter AD.: Urine proteomic profiling of pediatric nephrotic syndrome. Pediatr Nephrol. 2006 Sep;21(9):1257-65. Epub 2006 Jun 30.

**For laboratory research only, not for drug, diagnostic or other use.**