



A4-250-C025

Monoclonal Antibody to alpha-tubulin Alexa Fluor® 488 conjugated (0.025 mg)

| | |
|-----------------------------|---|
| Clone: | TU-01 |
| Isotype: | Mouse IgG1 |
| Specificity: | The antibody TU-01 recognizes the defined epitope (aa 65-97) on N-terminal structural domain of alpha-tubulin. |
| Immunogen: | Fraction of tubulin purified from porcine brain by two cycles of polymerization - depolymerization. |
| Species Reactivity: | Broad species reactivity |
| Preparation: | The purified antibody is conjugated with Alexa Fluor® 488 under optimum conditions. The conjugate is purified by size-exclusion chromatography. |
| Concentration: | 1.0 mg/ml |
| Storage Buffer: | Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4 |
| Storage / Stability: | Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user. |
| Usage: | The antibody is designed for Immunocytochemistry analysis. Suggested working dilution is 1:20-100. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator. |
| Expiration: | See vial label |
| Lot Number: | See vial label |

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****Background:**

The microtubules are intracellular dynamic polymers made up of evolutionarily conserved polymorphic alpha/beta-tubulin heterodimers and a large number of microtubule-associated proteins (MAPs). The microtubules consist of 13 protofilaments and have an outer diameter 25 nm. Microtubules have their intrinsic polarity; highly dynamic plus ends and less dynamic minus ends. Microtubules are required for vital processes in eukaryotic cells including mitosis, meiosis, maintenance of cell shape and intracellular transport. Microtubules are also necessary for movement of cells by means of flagella and cilia. In mammalian tissue culture cells microtubules have their minus ends anchored in microtubule organizing centers (MTOCs). The GTP (guanosin triphosphate) molecule is an essential for tubulin heterodimer to associate with other heterodimers to form microtubule. In vivo, microtubule dynamics vary considerably. Microtubule polymerization is reversible and a populations of microtubules in cells are on their minus ends either growing or shortening; this phenomenon is called dynamic instability of microtubules. On a practical level, microtubules can easily be stabilized by the addition of non-hydrolysable analogues of GTP (eg. GMPPCP) or more commonly by anti-cancer drugs such as Taxol. Taxol stabilizes microtubules at room temperature for many hours. Using limited proteolysis by enzymes both tubulin subunits can be divided into N-terminal and C-terminal structural domains. The alpha-tubulin (relative molecular weight around 50 kDa) is globular protein that exists in cells as part of soluble alpha/beta-tubulin dimer or it is polymerized into microtubules. In different species it is coded by multiple tubulin genes that form tubulin classes (in human 6 genes). Expressed tubulin genes are named tubulin isotypes. Some of the tubulin isotypes are expressed ubiquitously, while some have more restricted tissue expression. Alpha-tubulin is also subject of numerous post-translational modifications. Tubulin isotypes and their posttranslational modifications are responsible for multiple tubulin charge variants - tubulin isoforms. Heterogeneity of alpha-tubulin is concentrated in C-terminal structural domain.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Viklicky V, Draber P, Hasek J, Bartek J: Production and characterization of a monoclonal antitubulin antibody. *Cell Biol Int Rep.* 1982 Aug;6(8):725-31.
- *Draber P, Draberova E, Zicconi D, Sellitto C, Viklicky V, Cappuccinelli P: Heterogeneity of microtubules recognized by monoclonal antibodies to alpha-tubulin. *Eur J Cell Biol.* 1986 Jun;41(1):82-8.
- *Grimm M, Breitling F, Little M: Location of the epitope for the alpha-tubulin monoclonal antibody TU-O1. *Biochim Biophys Acta.* 1987 Jul 24;914(1):83-8.
- *Draber P, Draberova E, Linhartova I, Viklicky V: Differences in the exposure of C- and N-terminal tubulin domains in cytoplasmic microtubules detected with domain-specific monoclonal antibodies. *J Cell Sci.* 1989 Mar;92 (Pt 3):519-28.
- *Draber P, Draberova E, Viklicky V: Immunostaining of human spermatozoa with tubulin domain-specific monoclonal antibodies. Recognition of a unique beta tubulin epitope in the sperm head. *Histochemistry.* 1991;95(5):519-24.
- *Linhartova I, Draber P, Draberova E, Viklicky V: Immunological discrimination of beta-tubulin isoforms in developing mouse brain. Post-translational modification of non-class-III beta-tubulins. *Biochem J.* 1992 Dec 15;288 (Pt 3):919-24.
- *Nováková M, Dráberová E, Schürmann W, Czihak G, Viklický V, Dráber P: gamma-Tubulin redistribution in taxol-treated mitotic cells probed by monoclonal antibodies. *Cell Motil Cytoskeleton.* 1996;33(1):38-51.
- *Smertenko A, Blume Y, Viklicky V, Opatrny Z, Draber P: Post-translational modifications and multiple tubulin isoforms in *Nicotiana tabacum* L. cells. *Planta.* 1997;201(3):349-58.
- *Kukharsky V, Sulimenko V, Macůrek L, Sulimenko T, Dráberová E, Dráber P: Complexes of gamma-tubulin with nonreceptor protein tyrosine kinases Src and Fyn in differentiating P19 embryonal carcinoma cells. *Exp Cell Res.* 2004 Aug 1;298(1):218-28.
- *Lukas J, Mazna P, Valenta T, Doubravska L, Pospichalova V, Vojtechova M, Fafílek B, Ivanek R, Plachy J, Novak J, Korinek V: Dazap2 modulates transcription driven by the Wnt effector TCF-4. *Nucleic Acids Res.* 2009 Mar 20. [Epub ahead of print]
- *Smertenko A, Blume Y, Viklický V, Dráber P: Exposure of tubulin structural domains in *Nicotiana tabacum* microtubules probed by monoclonal antibodies. *Eur J Cell Biol.* 1997 Feb;72(2):104-12.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz