



11-382-C100

## Monoclonal Antibody to CD16 Purified Antibody (0.1 mg)

<b>Clone:</b>	MEM-168
<b>Isotype:</b>	Mouse IgM
<b>Specificity:</b>	The antibody MEM-168 reacts with CD16 antigen, a low affinity receptor for aggregated IgG (Fcγ <sub>3</sub> antigen). CD16 exists in two different isoforms: CD16a (Fcγ <sub>3A</sub> ; 50-65 kDa; expressed on NK-cells, monocytes and macrophages) and CD16b (Fcγ <sub>3B</sub> ; 48 kDa; mainly expressed on neutrophils).
<b>Immunogen:</b>	Human granulocytes
<b>Species Reactivity:</b>	Human, Non-Human Primates, Porcine
<b>Application:</b>	<b>Flow Cytometry</b>
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified from ascites by gel filtration and precipitation methods.
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not use after expiration date stamped on vial label. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	<b>CD16</b> (Fcγ <sub>3</sub> ) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human Fcγ <sub>3</sub> is expressed in two forms Fcγ <sub>3A</sub> and -B. Fcγ <sub>3A</sub> is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with Fcε <sub>1</sub> -γ subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell Fcγ <sub>3A</sub> is associated, moreover, with Fcε <sub>1</sub> -β subunit. Besides IgG, Fcγ <sub>3A</sub> can be triggered also by oligomeric IgE. Fcγ <sub>3B</sub> is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.
<b>References:</b>	*Gessner JE, Grussenmeyer T, Kolanus W, Schmidt RE: The human low affinity immunoglobulin G Fc receptor III-A and III-B genes. Molecular characterization of the promoter regions. J Biol Chem. 1995 Jan 20;270(3):1350-61. *Kocher M, Siegel ME, Edberg JC, Kimberly RP: Cross-linking of Fc gamma receptor IIa and Fc gamma receptor IIIb induces different proadhesive phenotypes on human neutrophils. J Immunol. 1997 Oct 15;159(8):3940-8. *Arase N, Arase H, Hirano S, Yokosuka T, Sakurai D, Saito T: IgE-mediated activation of NK cells through Fc gamma RIII. J Immunol. 2003 Mar 15;170(6):3054-8. *Drbal K, Moertelmaier M, Holzhauser C, Muhammad A, Fuertbauer E, Howorka S, Hinterberger M, Stockinger H, Schütz GJ: Single-molecule microscopy reveals heterogeneous dynamics of lipid raft components upon TCR engagement. Int Immunol. 2007 May;19(5):675-84.

**For laboratory research only, not for drug, diagnostic or other use.**