



Antibodies

PC-399-T025

Monoclonal Antibody to CD16 PerCP (25 tests)

Clone:	LNK16
Isotype:	Mouse IgG1
Specificity:	The antibody LNK16 reacts with CD16, a low affinity receptor for aggregated IgG (FcγR3 antigen). CD16 exists in two different isoforms: CD16a (FcγR3A; 50-65 kDa; expressed on NK-cells, monocytes and macrophages) and CD16b (FcγR3B; 48 kDa; mainly expressed on neutrophils). HLDA V; WS Code M MA069 HLDA V; WS Code NK NK50
Immunogen:	Normal human peripheral blood granulocytes
Species Reactivity:	Human, Non-Human Primates
Preparation:	The purified antibody is conjugated with Peridinin-chlorophyll-protein complex (PerCP) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.25 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD16 (FcγR3) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcγR3 is expressed in two forms – FcγR3-A and -B. FcγR3-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcεR1-γ subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcγR3-A is associated, moreover, with FcεR1-β subunit. Besides IgG, FcγR3-A can be triggered also by oligomeric IgE. FcγR3-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Gessner JE, Grussenmeyer T, Kolanus W, Schmidt RE: The human low affinity immunoglobulin G Fc receptor III-A and III-B genes. Molecular characterization of the promoter regions. *J Biol Chem.* 1995 Jan 20;270(3):1350-61.
- *Kocher M, Siegel ME, Edberg JC, Kimberly RP: Cross-linking of Fc gamma receptor IIa and Fc gamma receptor IIIb induces different proadhesive phenotypes on human neutrophils. *J Immunol.* 1997 Oct 15;159(8):3940-8.
- *Arase N, Arase H, Hirano S, Yokosuka T, Sakurai D, Saito T: IgE-mediated activation of NK cells through Fc gamma RIII. *J Immunol.* 2003 Mar 15;170(6):3054-8.
- *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- *Tamm A, Schmidt RE: The binding epitopes of human CD16 (Fc gamma RIII) monoclonal antibodies. Implications for ligand binding. *J Immunol.* 1996 Aug 15;157(4):1576-81.
- *Boyle JJ: Human macrophages kill human mesangial cells by Fas-L-induced apoptosis when triggered by antibody via CD16. *Clin Exp Immunol.* 2004 Sep;137(3):529-37.
- *Lin CW, Liu TY, Chen SU, Wang KT, Medeiros LJ, Hsu SM: CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. *Blood.* 2005 Nov 15;106(10):3567-74.
- *Mathison RD, Befus AD, Davison JS, Woodman RC: Modulation of neutrophil function by the tripeptide feG. *BMC Immunol.* 2003 Mar 4;4:3. Epub 2003 Mar 4.
- *Hovden AO, Karlsen M, Jonsson R, Aarstad HJ, Appel S: Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. *BMC Immunol.* 2011 Jan 5;12:2.

For laboratory research only, not for drug, diagnostic or other use.