



11-460-C025

Monoclonal Antibody to Vimentin Purified Antibody (0.025 mg)

Clone:	VI-10
Isotype:	Mouse IgM
Specificity:	The antibody VI-10 reacts with vimentin, a 57 kDa intermediate filament expressed in variety of mesenchymal and mesodermal cell types.
Regulatory Status:	RUO
Species Reactivity:	Human, Porcine, Mouse, Rat, Chicken, Other not tested
Application:	Immunoprecipitation Western Blotting Immunohistochemistry (paraffin sections) Recommended dilution: 5 µg/ml Positive tissue: skin fibroblast Immunocytochemistry Staining technique: RBL basophilic leukemia cell line: (a) Fix cells for 10 min in methanol at -20°C and for 6 min in acetone at -20°C; (b) Fix cells directly in methanol for 10 min at -20°C or in acetone for 10 min at -20°C. Incubation: 45 min RT Positive control: 3T3 mouse Swiss albino fibroblast cell line RBL rat basophilic cell line
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by precipitation and chromatography
Concentration:	1 mg/ml
Storage Buffer:	Tris buffered saline (TBS) with 15 mM sodium azide, approx. pH 8.0
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****Background:**

Vimentin (57 kDa) is the most ubiquitous intermediate filament protein and the first to be expressed during cell differentiation. All primitive cell types express vimentin but in most non-mesenchymal cells it is replaced by other intermediate filament proteins during differentiation. Vimentin is expressed in a wide variety of mesenchymal cell types - fibroblasts, endothelial cells etc., and in a number of other cell types derived from mesoderm, e.g., mesothelium and ovarian granulosa cells. In non-vascular smooth muscle cells and striated muscle, vimentin is often replaced by desmin, however, during regeneration, vimentin is reexpressed. Cells of the lympho-haemopoietic system (lymphocytes, macrophages etc.) also express vimentin, sometimes in scarce amounts. Vimentin is also found in mesoderm derived epithelia, e.g. kidney (Bowman capsule), endometrium and ovary (surface epithelium), in myoepithelial cells (breast, salivary and sweat glands), and in thyroid gland epithelium. In these cell types, as in mesothelial cells, vimentin is coexpressed with cytokeratin.

Furthermore, vimentin is detected in many cells from the neural crest. Particularly melanocytes express abundant vimentin. In glial cells vimentin is coexpressed with Glial Fibrillary Acidic Protein (GFAP).

Vimentin is present in many different neoplasms but is particularly expressed in those originated from mesenchymal cells. Sarcomas e.g., fibrosarcoma, malignant fibrous histiocytoma, angiosarcoma, and leiomyosarcoma, as well as lymphomas, malignant melanoma and schwannoma, are virtually always vimentin positive. Mesoderm derived carcinomas like renal cell carcinoma, adrenal cortical carcinoma and adenocarcinomas from endometrium and ovary usually express vimentin. Also thyroid carcinomas are vimentin positive. Any low differentiated carcinoma may express some vimentin.

Vimentin is frequently included in the so-called primary panel (together with CD45, cytokeratin, and S-100 protein). Intense staining reaction for vimentin without coexpression of other intermediate filament proteins is strongly suggestive of a mesenchymal tumour or malignant melanoma.

References:

*Draberova E, Zikova M, Draber P: Monoclonal antibody VI-10 specific for vimentin. *Folia Biol (Praha)*. 1999;45(1):35-6.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's terms and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.