

1A-458-C100

## Mouse IgG2a Isotype Control Allophycocyanin (APC) conjugated (0.1 mg)

Clone: PPV-04

**Isotype:** Mouse IgG2a

Specificity: This mouse IgG2a monoclonal antibody (clone PPV-04) reacts with undefined

epitope on a plant pathogen.

Regulatory Status: RUO

Negative Species: Human, Porcine, Mouse, Rat, Other species Not tested (expected negative)

**Preparation:** The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under

optimum conditions. The conjugate is purified by size-exclusion chromatography.

Concentration: 0.1 mg/ml

Storage Buffer: The reagent is provided in stabilizing phosphate buffered saline (PBS) solution

containing 15mM sodium azide.

Storage / Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not

use after expiration date stamped on vial label.

**Usage:** The reagent is intended as isotype control for flow cytometry analysis to establish

the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this isotype control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above 10 µg per ml of incubation mixture), change the conditions of

your experiment to reduce the background.

Expiration: See vial label

Lot Number: See vial label

**Background:** The specificity of staining by monoclonal antibodies to target antigens should be

verified by establishing the amount of non-specific antibody binding. Especially at higher concentration (more than 15  $\mu$ g/ml) the antibody staining usually has consignable background. To this end a non-reactive immunoglobulin of the same isotype is included as a negative control for each specific monoclonal antibody used in a particular immunoassay. The monoclonal antibody PPV-04, generated against a plant pathogen, does not cross-react with other species, and hence all the background that could be observed when working with this antibody would be a result of general nonspecific interactions between an mouse IgG2a molecule and the respective sample under the particular conditions. This shall help the customer to set up the experimental conditions so that the nonspecific binding of any

antibody is abolished.

References: Unpublished.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.