

PC-531-T025

Monoclonal Antibody to CD86 PerCP (25 tests)

Clone:	BU63
Isotype:	Mouse IgG1
Specificity:	<p>The antibody BU63 reacts with CD86 (B7-2), a 70 kDa type I transmembrane glycoprotein of immunoglobulin supergene family, expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes.</p> <p>HLDA V; WS Code BP BP072 HLDA V; WS Code A A109 HLDA VI; WS Code BP 95 HLDA VI; WS Code B CD86.9</p>
Immunogen:	B-lymphoblastoid cell line ARH 77
Species Reactivity:	Human, Other not determined
Preparation:	The purified antibody is conjugated with Peridinin-chlorophyll-protein complex (PerCP) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	<p>Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.</p> <p>Do not use after expiration date stamped on vial label.</p> <p>Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.</p>
Usage:	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.</p> <p>The content of a vial (0.25 ml) is sufficient for 25 tests.</p>
Expiration:	See vial label
Lot Number:	See vial label
Background:	<p>CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, the both molecules can also substitute for each other in this process. The question what are the differences in CD80 and CD86 competency has not been fully elucidated yet; there are still conflicts in results about their respective roles in initiation or sustaining of the T cell immune response.</p>

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Vasilevko V, Ghochikyan A, Holterman MJ, Agadjanyan M.G.: CD80 (B7-1) and CD86 (B7-2) are functionally equivalent in the initiation and maintenance of CD4+ T-cell proliferation after activation with suboptimal doses of PHA. *DNA Cell Biol.* 2002 Mar;21(3):137-49.
- *Yadav D, Judkowski V, Flodstrom-Tullberg M, Sterling L, Redmond WL, Sherman L, Sarvetnick N.: B7-2 (CD86) controls the priming of autoreactive CD4 T cell response against pancreatic islets. *J Immunol.* 2004 Sep 15;173(6):3631-9.
- *Thomas IJ, Petrich de Marquesini LG, Ravanan R, Smith RM, Guerder S, Flavell RA, Wraith DC, Wen L, Wong F.S.: CD86 has sustained costimulatory effects on CD8 T cells. *J Immunol.* 2007 Nov 1;179(9):5936-46.
- *Eri R, Kodumudi KN, Summerlin DJ, Srinivasan M.: Suppression of colon inflammation by CD80 blockade: Evaluation in two murine models of inflammatory bowel disease. *Inflamm Bowel Dis.* 2008 Jan 9
- *Engel P, Gribben JG, Freeman GJ, Zhou LJ, Nozawa Y, Abe M, Nadler LM, Wakasa H, Tedder T.F.: The B7-2 (B70) costimulatory molecule expressed by monocytes and activated B lymphocytes is the CD86 differentiation antigen. *Blood.* 1994 Sep 1;84(5):1402-7.
- *Caux C, Vanbervliet B, Massacrier C, Azuma M, Okumura K, Lanier LL, Banchereau J.: B70/B7-2 is identical to CD86 and is the major functional ligand for CD28 expressed on human dendritic cells. *J Exp Med.* 1994 Nov 1;180(5):1841-7.
- *Mauri D, Wyss-Coray T, Gallati H, Pichler W.J.: Antigen-presenting T cells induce the development of cytotoxic CD4+ T cells. I. Involvement of the CD80-CD28 adhesion molecules. *J Immunol.* 1995 Jul 1;155(1):118-27.
- *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- *Giguère JF, Bounou S, Paquette JS, Madrenas J, Tremblay M.J.: Insertion of host-derived costimulatory molecules CD80 (B7.1) and CD86 (B7.2) into human immunodeficiency virus type 1 affects the virus life cycle. *J Virol.* 2004 Jun;78(12):6222-32.

For laboratory research only, not for drug, diagnostic or other use.