

A4-588-T025

Monoclonal Antibody to CD324 / E-Cadherin Alexa Fluor® 488 conjugated (25 tests)

Clone:	67A4
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody 67A4 recognizes CD324 / E-cadherin, an approximately 100 kDa epithelial cell adhesion molecule, whose detection is important for determination of invasive potential of epithelial neoplasms. HLDA VIII
Immunogen:	T-47D cells
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Alexa Fluor® 488 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD324 / E-cadherin is an epithelial cell surface molecule, which provides calcium-dependent homophilic interactions with E-cadherin of another cell. These interactions take part in morphogenetic programs controlling the maintenance of the structural and functional integrity of epithelia and affect invasive potential of epithelial neoplasms. CD324 / E-cadherin is implicated in cell growth and differentiation, cell recognition, and sorting during developmental morphogenesis, as well as in aggregation-dependent cell survival. CD324 / E-cadherin-mediated cell adhesion system is highly regulated from inside the cell by a number of intracellular signaling pathways.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Takeichi M: Cadherin cell adhesion receptors as a morphogenetic regulator. *Science*. 1991 Mar 22;251(5000):1451-5.
- *Pece S, Chiariello M, Murga C, Gutkind JS: Activation of the protein kinase Akt/PKB by the formation of E-cadherin-mediated cell-cell junctions. Evidence for the association of phosphatidylinositol 3-kinase with the E-cadherin adhesion complex. *J Biol Chem*. 1999 Jul 2;274(27):19347-51.
- *Pece S, Gutkind JS: Signaling from E-cadherins to the MAPK pathway by the recruitment and activation of epidermal growth factor receptors upon cell-cell contact formation. *J Biol Chem*. 2000 Dec 29;275(52):41227-33.
- *Armeanu S, Bühring HJ, Reuss-Borst M, Müller CA, Klein G: E-cadherin is functionally involved in the maturation of the erythroid lineage. *J Cell Biol*. 1995 Oct;131(1):243-9.
- *Bühring HJ, Müller T, Herbst R, Cole S, Rappold I, Schuller W, Zhu X, Fritzscht U, Faul C, Armeanu S, Ullrich A, Klein G, Schmidt H: The adhesion molecule E-cadherin and a surface antigen recognized by the antibody 9C4 are selectively expressed on erythroid cells of defined maturational stages. *Leukemia*. 1996 Jan;10(1):106-16.
- *Novak N, Kraft S, Haberstock J, Geiger E, Allam P, Bieber T: A reducing microenvironment leads to the generation of FcepsilonRIhigh inflammatory dendritic epidermal cells (IDEC). *J Invest Dermatol*. 2002 Oct;119(4):842-9.
- *Servet-Delprat C, Vidalain PO, Bausinger H, Manié S, Le Deist F, Azocar O, Hanau D, Fischer A, Rabourdin-Combe C: Measles virus induces abnormal differentiation of CD40 ligand-activated human dendritic cells. *J Immunol*. 2000 Feb 15;164(4):1753-60.
- *Furio L, Guezennec A, Ducarre B, Guesnet J, Peguet-Navarro J: Differential effects of allergens and irritants on early differentiating monocyte-derived dendritic cells. *Eur J Dermatol*. 2008 Mar-Apr;18(2):141-7.
- *Robertson H, Ali S, McDonnell BJ, Burt AD, Kirby JA: Chronic renal allograft dysfunction: the role of T cell-mediated tubular epithelial to mesenchymal cell transition. *J Am Soc Nephrol*. 2004 Feb;15(2):390-7.
- *Kutlesa S, Wessels JT, Speiser A, Steiert I, Müller CA, Klein G: E-cadherin-mediated interactions of thymic epithelial cells with CD103+ thymocytes lead to enhanced thymocyte cell proliferation. *J Cell Sci*. 2002 Dec 1;115(Pt 23):4505-15.
- *Caberg JH, Hubert PM, Begon DY, Herfs MF, Roncarati PJ, Boniver JJ, Delvenne PO: Silencing of E7 oncogene restores functional E-cadherin expression in human papillomavirus 16-transformed keratinocytes. *Carcinogenesis*. 2008 Jul;29(7):1441-7.
- *Lin JC, Liao SK, Lee EH, Hung MS, Sayion Y, Chen HC, Kang CC, Huang LS, Cherng JM. Molecular events associated with epithelial to mesenchymal transition of nasopharyngeal carcinoma cells in the absence of Epstein-Barr virus genome. *J Biomed Sci*. 2009 Nov 24;16:105.
- *And many other.

This product is provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Exbio Praha, a.s., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including use in flow cytometry that does not utilize a bead based array, but excluding use in combination with microarrays or High Content Screening. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz