



1F-621-C100

Monoclonal Antibody to MHC Class II (mouse) Fluorescein (FITC) conjugated (0.1 mg)

Clone:	M5/114
Isotype:	Rat IgG2b
Specificity:	The rat monoclonal antibody M5/114 reacts with murine MHC class II glycoproteins. It recognizes a shared determinant on I-Ab, I-Ad, I-Aq, and I-Ed, I-Ek alloantigens, but it does not react with I-Af, I-Ak, I-As. This antibody can inhibit I-A-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not H-2f, H-2k, H-2s haplotypes.
Immunogen:	Activated C57BL/6 mouse spleen cells
Species Reactivity:	Mouse
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.
Usage:	The reagent is designed for Flow Cytometry analysis. Suggested working concentration is 4 µg/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	MHC (major histocompatibility complex) class II molecules are transmembrane glycoproteins expressed on the surface of professional antigen-presenting cells, such as macrophages, dendritic cells and B cells. Before their exposition on the cell surface, the MHC class II molecules react with endocytosed exogenous antigens, which are then presented to the T cells. The antigen-binding groove between MHC class II alpha and beta chain is open at both ends and is 15-24 amino acid residues long.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Bhattacharya A, Dorf ME, Springer TA: A shared alloantigenic determinant on Ia antigens encoded by the I-A and I-E subregions: evidence for I region gene duplication. *J Immunol.* 1981 Dec;127(6):2488-95.
- *Viville S, Neefjes J, Lotteau V, Dierich A, Lemeur M, Ploegh H, Benoist C, Mathis D: Mice lacking the MHC class II-associated invariant chain. *Cell.* 1993 Feb 26;72(4):635-48.
- *De Souza Leao S, Lang T, Prina E, Hellio R, Antoine JC: Intracellular *Leishmania amazonensis* amastigotes internalize and degrade MHC class II molecules of their host cells. *J Cell Sci.* 1995 Oct;108 (Pt 10):3219-31.
- *Kleijmeer M, Ramm G, Schuurhuis D, Griffith J, Rescigno M, Ricciardi-Castagnoli P, Rudensky AY, Ossendorp F, Melief CJ, Stoorvogel W, Geuze HJ: Reorganization of multivesicular bodies regulates MHC class II antigen presentation by dendritic cells. *J Cell Biol.* 2001 Oct 1;155(1):53-63.
- *Beers C, Burich A, Kleijmeer MJ, Griffith JM, Wong P, Rudensky AY: Cathepsin S controls MHC class II-mediated antigen presentation by epithelial cells in vivo. *J Immunol.* 2005 Feb 1;174(3):1205-12.
- *Kuwano Y, Prazma CM, Yazawa N, Watanabe R, Ishiura N, Kumanogoh A, Okochi H, Tamaki K, Fujimoto M, Tedder TF: CD83 influences cell-surface MHC class II expression on B cells and other antigen-presenting cells. *Int Immunol.* 2007 Aug;19(8):977-92.
- *Zang W, Kalache S, Lin M, Schroppel B, Murphy B: MHC Class II-mediated apoptosis by a nonpolymorphic MHC Class II peptide proceeds by activation of protein kinase C. *J Am Soc Nephrol.* 2005 Dec;16(12):3661-8.
- *Clausen BE, Waldburger JM, Schwenk F, Barras E, Mach B, Rajewsky K, Förster I, Reith W: Residual MHC class II expression on mature dendritic cells and activated B cells in RFX5-deficient mice. *Immunity.* 1998 Feb;8(2):143-55.
- *And many other.

For laboratory research only, not for drug, diagnostic or other use.