

1F-622-C500

Monoclonal Antibody to CD11a (mouse) Fluorescein (FITC) conjugated (0.5 mg)

Clone:	M17/4
Isotype:	Rat IgG2a
Specificity:	The rat monoclonal antibody M17/4 reacts with CD11a (alpha-subunit of murine LFA-1), a 180 kDa type I transmembrane glycoprotein expressed on B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils.
Immunogen:	C57BL/6 mouse splenic secondary cytotoxic T lymphocytes
Species Reactivity:	Mouse
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.
Usage:	The reagent is designed for Flow Cytometry analysis. Suggested working concentration is 3 µg/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity conformation. Cell stimulation by chemokines or other signals leads to induction the high-affinity conformation, which supports tight binding of LFA-1 to its ligands, the intercellular adhesion molecules ICAM-1, -2, -3. LFA-1 is thus involved in interaction of various immune cells and in their tissue-specific settlement, but participates also in control of cell differentiation and proliferation and of T-cell effector functions.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Sanchez-Madrid F, Davignon D, Martz E, Springer TA: Antigens involved in mouse cytolytic T-lymphocyte (CTL)-mediated killing: functional screening and topographic relationship. *Cell Immunol.* 1982 Oct;73(1):1-11.
- *Kuhlman P, Moy VT, Lollo BA, Brian AA: The accessory function of murine intercellular adhesion molecule-1 in T lymphocyte activation. Contributions of adhesion and co-activation. *J Immunol.* 1991 Mar 15;146(6):1773-82.
- *Angelov GS, Guillaume P, Cebecauer M, Bosshard G, Dojcinovic D, Baumgaertner P, Luescher IF: Soluble MHC-peptide complexes containing long rigid linkers abolish CTL-mediated cytotoxicity. *J Immunol.* 2006 Mar 15;176(6):3356-65.
- *Smith SS, Barnum SR: Differential expression of beta 2-integrins and cytokine production between gammadelta and alphabeta T cells in experimental autoimmune encephalomyelitis. *J Leukoc Biol.* 2008 Jan;83(1):71-9.
- *Yung R, Powers D, Johnson K, Amento E, Carr D, Laing T, Yang J, Chang S, Hemati N, Richardson B: Mechanisms of drug-induced lupus. II. T cells overexpressing lymphocyte function-associated antigen 1 become autoreactive and cause a lupuslike disease in syngeneic mice. *J Clin Invest.* 1996 Jun 15;97(12):2866-71.
- *Bellingan GJ, Xu P, Cooksley H, Cauldwell H, Shock A, Bottoms S, Haslett C, Mutsaers SE, Laurent GJ: Adhesion molecule-dependent mechanisms regulate the rate of macrophage clearance during the resolution of peritoneal inflammation. *J Exp Med.* 2002 Dec 2;196(11):1515-21.
- *Beyer M, Wang H, Peters N, Doths S, Koerner-Rettberg C, Openshaw PJ, Schwarze J: The beta2 integrin CD11c distinguishes a subset of cytotoxic pulmonary T cells with potent antiviral effects in vitro and in vivo. *Respir Res.* 2005 Jul 12;6:70.
- *Koseki S, Miura S, Fujimori H, Hokari R, Komoto S, Hara Y, Ogino T, Nagata H, Goto M, Hachimura S, Kaminogawa S, Ishii H: In situ demonstration of intraepithelial lymphocyte adhesion to villus microvessels of the small intestine. *Int Immunol.* 2001 Sep;13(9):1165-74.
- *Papayannopoulou T, Priestley GV, Nakamoto B, Zafiroopoulos V, Scott LM, Harlan JM: Synergistic mobilization of hemopoietic progenitor cells using concurrent beta1 and beta2 integrin blockade or beta2-deficient mice. *Blood.* 2001 Mar 1;97(5):1282-8.
- *And many other.

For laboratory research only, not for drug, diagnostic or other use.