

Technical Data Sheet

Product **Anti-Hu CD64 Purified Low Endotoxin**

Cat. Number/Size **12-644-C100 0.1 mg**

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Antigen	CD64
Clone	10.1
Format	Purified Low Endotoxin
Reactivity	Human, Non-human primates
Application	FC (QC tested), IP, WB, IHC(P), IHC(F), ICC, FUNC
Application details	Functional application: Blocking of IgG binding to the FcγR1. Flow cytometry: Recommended dilution: 1-4 µg/ml. Immunohistochemistry (paraffin sections and frozen sections): There can occur problems with paraformaldehyde fixation.
Isotype	Mouse IgG1 kappa
Specificity	The mouse monoclonal antibody 10.1 recognizes an extracellular epitope on CD64/FcγRI, a 72 kDa single chain type I glycoprotein, that is expressed on monocytes/macrophages, dendritic cells, and activated granulocytes. The epitope is sensitive to formalin fixation.
Other names	FcRI, IGFR1, FcγR1A
Workshop	HLDA III: WS Code M-250
Immunogen	Rheumatoid synovial fluid cells and fibronectin purified human monocytes
Entrez Gene ID	2209
Gene name	FCGR1A
NCBI Full Gene Name	Fc fragment of IgG receptor Ia
UniProt ID	P12314

Concentration	1 mg/ml
Preparation	Purified by protein-A affinity chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.