



	Technical Data Sheet	
Product	<b>Anti-Hu CD25 Purified</b>	
Cat. Number/Size	<b>11-218-C025</b>	<b>0.025 mg</b>
	<b>11-218-C100</b>	<b>0.1 mg</b>
	<b>For Research Use Only.</b>	
	<b>Not for use in diagnostic or therapeutic procedures.</b>	
Antigen	CD25	
Clone	MEM-181	
Format	Purified	
Reactivity	Human	
Application	FC (QC tested)	
Application details	Flow cytometry: Recommended dilution: 1-5 µg/ml.	
Isotype	Mouse IgG1 kappa	
Specificity	The antibody MEM-181 reacts with an extracellular epitope of CD25 (Interleukin-2 receptor alpha chain), a 55 kDa type I transmembrane glycoprotein expressed on activated B and T lymphocytes, activated monocytes/macrophages and on CD4+ T lymphocytes (T regulatory cells); it is lost on resting B and T lymphocytes.	
Other names	IL-2R α chain, IL-2-RA, IL2-RA, p55, TAC antigen, IDDM10, TCGFR	
Workshop	HLDA VI: WS Code NL N-L024	
Immunogen	PHA-activated peripheral blood leucocytes	
Entrez Gene ID	3559	
Gene name	IL2RA	
NCBI Full Gene Name	interleukin 2 receptor subunit alpha	
UniProt ID	P01589	
Concentration	1 mg/ml	
Preparation	Purified by protein-A affinity chromatography.	
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Storage and handling	Store at 2-8°C. Do not freeze.	
Images and References	www.exbio.cz	

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.