



	Technical Data Sheet	
Product	Anti-Hu CD9 Purified Azide Free	
Cat. Number/Size	10-208-C025	0.025 mg
	10-208-C100	0.1 mg
	For Research Use Only.	
	Not for use in diagnostic or therapeutic procedures.	
Antigen	CD9	
Clone	MEM-61	
Format	Purified Azide Free	
Reactivity	Human	
Application	FC (QC tested), WB, IHC-P, MC-CyTOF, FUNC	
Application details	Functional application: The antibody MEM-61 induces FcγR-dependent platelet aggregation. Flow cytometry: Recommended dilution: 1-4 µg/ml. Western blotting: Recommended dilution: 2-4 µg/ml, non-reducing conditions. Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/ml, positive tissue: prostate.	
Isotype	Mouse IgG1	
Specificity	The antibody MEM-61 recognizes an epitope on second extracellular domain (EC2) of CD9 antigen, a 24 kDa transmembrane protein expressed on platelets, monocytes, pre-B lymphocytes, granulocytes and activated T lymphocytes.	
Other names	MIC3, MRP-1, BTCC-1, DRAP-27, TSPAN29, BA2	
Workshop	HLDA VI: WS Code P P-15	
Immunogen	Pre-B cell line NALM-6.	
Entrez Gene ID	928	
Gene name	CD9	
NCBI Full Gene Name	CD9 molecule	
UniProt ID	P21926	
Concentration	1 mg/ml	
Preparation	Purified by protein-A affinity chromatography.	
Formulation	Phosphate buffered saline (PBS), pH 7.4	
Storage and handling	Store at 2-8°C. Do not freeze.	
Images and References	www.exbio.cz	

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.