

Technical Data Sheet

Product	Anti-Hu CD54 Purified	
Cat. Number/Size	11-229-C025	0.025 mg
	11-229-C100	0.1 mg
	For Research Use Only.	
	Not for use in diagnostic or therapeutic procedures.	

Antigen	CD54
Clone	MEM-112
Format	Purified
Reactivity	Human
Application	FC (QC tested), IP
Application details	Flow cytometry: Recommended dilution: 0.5-4 µg/ml.
Isotype	Mouse IgG1
Specificity	The antibody MEM-112 reacts with an extracellular epitope of CD54 (ICAM-1), a 85-110 kDa type I transmembrane glycoprotein (receptor for rhinovirus). The expression of CD54 is upregulated by activation; it is expressed on activated endothelial cells, T lymphocytes, B lymphocytes, monocytes, macrophages, granulocytes and dendritic cells.
Other names	ICAM-1, BB2, P3.58
Workshop	HLDA VI: WS Code AS A050
Immunogen	Raji Burkitt's lymphoma cell line Raji
Entrez Gene ID	3383
Gene name	ICAM1
NCBI Full Gene Name	intercellular adhesion molecule 1
UniProt ID	P05362

Concentration	1 mg/ml
Preparation	Purified by protein-A affinity chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.