

**Technical Data Sheet** 

Product Anti-HLA-G Purified

Cat. Number/Size **11-499-C025 0.025 mg** 

11-499-C100 0.1 mg

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Antigen HLA-G
Clone 4H84
Format Purified
Reactivity Human

Application WB, IP, IHC-P, ICC, ELISA, IHC-F, FC

Application details Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/ml.

ELISA: The antibody 4H84 has been tested as the capture antibody in a sandwich ELISA for analysis of heavy

chains of all HLA-G isoforms in combination with antibody MEM-G/1 (cat. no. 1B-291-C100).

Flow cytometry: Recommended dilution: 2-5 µg/ml. Intracellular staining or surface staining after acid

treatment (Polakova K. et al. Molecular immunology 1993; 1223-30).

Western blotting: Recommended dilution 1-2 µg/ml.

Isotype Mouse IgG1

Specificity The mouse monoclonal antibody 4H84 recognizes an extracellular epitope of HLA-G molecule (39 kDa).

HLA-G belongs to the MHC Class I molecules (MHC Class Ib; nonclassical) and it is expressed on the surface

of trophoblast cells.

Immunogen amino acids 61-83 of HLA-G of human origin

Entrez Gene ID 3135
Gene name HLA-G

NCBI Full Gene Name major histocompatibility complex, class I, G

UniProt ID P17693

Concentration 1 mg/ml

Preparation Purified by protein-A affinity chromatography.

Formulation Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage and handling Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.

This product is for research and in vitro experimental use only. It is not to be used for any other commercial purpose. Use of this product to produce products for sale or for therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for commercial purposes, contact The Regents of the University of California.

Revision date: 2025-01-02