

Technical Data Sheet

Product

Anti-Transferrin Purified

Cat. Number/Size

11-258-C025

0.025 mg

11-258-C100

0.1 mg

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Antigen	Transferrin
Clone	HTF-14
Format	Purified
Reactivity	Human, Pig, Rabbit
Negative species	Horse, Dog, Sheep, Cow
Application	WB, IHC(P), ICC, ELISA, RIA
Application details	Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/ml; positive tissue: placenta. Western blotting: non-reducing conditions, recommended dilution: 1-2 µg/ml.
Isotype	Mouse IgG1
Specificity	The antibody HTF-14 recognizes an epitope located in the N-terminal domain of human serum transferrin, a 77 kDa single polypeptide chain glycoprotein (member of the iron binding family of proteins). It is synthesised in the liver and consists of two domains each having a high affinity reversible binding site for Fe ³⁺ .
Other names	TFQTL1, PRO1557, PRO2086
Immunogen	Purified porcine transferrin.
Entrez Gene ID	7018
Gene name	TF
NCBI Full Gene Name	transferrin
UniProt ID	P02787

Concentration	1 mg/ml
Preparation	Purified by protein-A affinity chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.