



	Technical Data Sheet	
Product	Anti-Hu CD99R Purified	
Cat. Number/Size	11-236-C025	0.025 mg
	11-236-C100	0.1 mg
	For Research Use Only.	
	Not for use in diagnostic or therapeutic procedures.	
Antigen	CD99R	
Clone	MEM-131	
Format	Purified	
Reactivity	Human	
Application	FC (QC tested), IP, WB, IHC-P	
Application details	Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/ml; positive tissue: thymic lymphocytes. Flow cytometry: Recommended dilution: 1 µg/ml. Western blotting: Non-reducing conditions.	
Isotype	Mouse IgM	
Specificity	The antibody MEM-131 reacts with CD99R, an extracellular epitope restricted to a subset of CD99 molecule expressed on myeloid cells, NK cells and T lymphocytes.	
Other names	MIC2, HBA71, blood group Xg	
Workshop	HLDA V: WS Code AS S020; HLDA V: WS Code T T-E2.02; HLDA V: WS Code T T-017	
Immunogen	HPB-ALL human peripheral blood leukemia T-cell line	
Concentration	1 mg/ml	
Preparation	Purified by sequential steps of physicochemical fractionation (differential precipitation and solid-phase chromatography methods).	
Formulation	Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	
Storage and handling	Store at 2-8°C. Do not freeze.	
Images and References	www.exbio.cz	

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.