



	Technical Data Sheet	
Product	Anti-Hu CD77 FITC	
Cat. Number/Size	1F-193-T100	100 tests
	For Research Use Only.	
	Not for use in diagnostic or therapeutic procedures.	
Antigen	CD77	
Clone	38.13	
Format	FITC	
Reactivity	Human	
Application	FC (QC tested)	
Application details	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.	
Excitation laser	blue (488 nm)	
Isotype	Lewis Rat IgM	
Specificity	The rat monoclonal antibody 38.13 recognizes CD77 (globotriaosylceramide Gb3), a neutral glycosphingolipid expressed mainly on the surface of B cell populations, such as germinal center B cells and Burkitt's lymphoma cells.	
Other names	globotriaosylceramide Gb3, Gal-alpha1-4Gal-beta1-4Glc-beta1-Cer, P blood group	
Workshop	HLDA IV	
Immunogen	Daudi cell line (Burkitt's lymphoma)	
Preparation	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.	
Formulation	Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	
Storage and handling	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.	
Images and References	www.exbio.cz	

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.