

Technical Data Sheet

Product	Anti-Hu CD56 Biotin	
Cat. Number/Size	1B-231-C025	0.025 mg
	1B-231-C100	0.1 mg
	For Research Use Only.	
	Not for use in diagnostic or therapeutic procedures.	

Antigen	CD56
Clone	MEM-188
Format	Biotin
Reactivity	Non-human primates, Human
Application	FC (QC tested), IP, IHC-P
Application details	Flow cytometry: Recommended dilution: 2-5 µg/ml.
Isotype	Mouse IgG2a
Specificity	The antibody MEM-188 reacts with an extracellular epitope on a 180 kDa isoform of CD56 (NCAM) expressed in leukocytes. It has been suggested that the antibody MEM-188 could react with rhesus monkey lymphocytes. Reactivity with other NCAM isoforms has not been tested.
Other names	NCAM1, MSK39
Workshop	HLDA VI: WS code A055; HLDA VI: WS Code NK26; HLDA VII: WS code 70077
Immunogen	KG-1 human acute myelogenous leukemia cell line
Entrez Gene ID	4684
Gene name	NCAM1
NCBI Full Gene Name	neural cell adhesion molecule 1
UniProt ID	P13591

Concentration	1 mg/ml
Preparation	Purified antibody is conjugated with biotin LC-NHS ester under optimum conditions and unconjugated antibody and free biotin are removed by size-exclusion chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.