



	Technical Data Sheet	
Product	<b>Anti-Hu CD8 Purified</b>	
Cat. Number/Size	<b>11-349-C025</b>	<b>0.025 mg</b>
	<b>11-349-C100</b>	<b>0.1 mg</b>
	<b>For Research Use Only.</b>	
	<b>Not for use in diagnostic or therapeutic procedures.</b>	
Antigen	CD8	
Clone	MEM-87	
Format	Purified	
Reactivity	Human	
Application	FC (QC tested), IP	
Application details	Flow cytometry: Recommended dilution: 2 µg/ml. Immunoprecipitation: Excellent for immunoisolation of CD8 positive T cells.	
Isotype	Mouse IgG1	
Specificity	The antibody MEM-87 recognizes an extracellular epitope of CD8, a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. CD8 is a disulfide-linked dimer and exists as a CD8 alpha/alpha homodimer or CD8 alpha/beta heterodimer (each monomer approx. 32-34 kDa).	
Other names	p32, LEU2	
Immunogen	Leukocytes of a patient suffering from LGL Type Leukemia.	
Concentration	1 mg/ml	
Preparation	Purified by protein-A affinity chromatography.	
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Storage and handling	Store at 2-8°C. Do not freeze.	
Images and References	www.exbio.cz	

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.