

Technical Data Sheet

Product	Anti-Hu CD55 Purified	
Cat. Number/Size	11-230-C025	0.025 mg
	11-230-C100	0.1 mg
	For Research Use Only.	
	Not for use in diagnostic or therapeutic procedures.	

Antigen	CD55
Clone	MEM-118
Format	Purified
Reactivity	Human, Non-human primates
Application	FC (QC tested), IP, IHC(P), WB
Application details	Flow cytometry: Recommended dilution: 10 µg/ml. Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/ml.
Isotype	Mouse IgM
Specificity	The antibody MEM-118 recognizes an epitope in SCR4 domain of CD55 (Decay accelerating factor, DAF), a 60-70 kDa glycosylphosphatidylinositol (GPI)-anchored single chain extracellular glycoprotein. CD55 is widely expressed on hematopoietic and on many non-hematopoietic cells; it is weakly present on NK cells.
Other names	DAF, CROM, CHAPLE
Workshop	HLDA V: WS Code AS S016
Immunogen	HPB-ALL human T cell line
Entrez Gene ID	1604
Gene name	CD55
NCBI Full Gene Name	CD55 molecule (Cromer blood group)
UniProt ID	P08174

Concentration	1 mg/ml
Preparation	Purified by sequential steps of physicochemical fractionation (differential precipitation and solid-phase chromatography methods).
Formulation	Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.