

Technical Data Sheet

Product	Anti-Hu CD142 Purified	
Cat. Number/Size	11-130-C025	0.025 mg
	11-130-C100	0.1 mg
	For Research Use Only.	
	Not for use in diagnostic or therapeutic procedures.	

Antigen	CD142
Clone	HTF-1
Format	Purified
Reactivity	Human
Application	FC (QC tested), WB, ICC, IHC(P)
Application details	Flow cytometry: Recommended dilution: 3-12 µg/ml. Western blotting: Recommended dilution: 1-2 µg/ml.
Isotype	Mouse IgG1 kappa
Specificity	The mouse monoclonal antibody HTF-1, also known as HTF1-7B8, recognizes an extracellular epitope of CD142 (tissue factor, coagulation factor III), a type I glycoprotein expressed on endothelial cells, monocytes, macrophages, and platelets upon induction by inflammatory mediators, and expressed constitutively by some tumors, the vasculature, placenta, kidney, and central nervous system.
Other names	F3, tissue factor, tissue thromboplastin, coagulation factor III, TF, TFA
Workshop	HLDA VI: WS Code E016
Immunogen	Human brain tissue factor (CD142)
Entrez Gene ID	2152
Gene name	F3
NCBI Full Gene Name	coagulation factor III, tissue factor
UniProt ID	P13726

Concentration	1 mg/ml
Preparation	Purified by protein-A affinity chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.