

Technical Data Sheet

Product

Anti-Hu CD16 Purified

Cat. Number/Size

11-214-C025

0.025 mg

11-214-C100

0.1 mg

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Antigen	CD16
Clone	MEM-154
Format	Purified
Reactivity	Human
Application	FC (QC tested), IP, WB
Application details	Flow cytometry: Recommended dilution: 5-10 µg/ml; positive control: PBL (peripheral blood lymphocytes). The antibody MEM-154 does not react with CD16a present on NK cells in many subjects. Western blotting: Non-reducing conditions.
Isotype	Mouse IgG1
Specificity	The antibody MEM-154 reacts with an extracellular epitope on CD16 antigen that is residing in proximity to FG loop (probably BC or C'E loop). CD16 is a low affinity receptor for aggregated IgG (Fcγ ₃ antigen). The antibody MEM-154 reacts with CD16 ⁺ granulocytes, and it can be used for mapping CD16-158V/F polymorphism on NK cells, as it requires presence of V at amino acid 158.
Other names	Fcγ ₃ , IGFR3, FCRIII
Workshop	HLDA V: WS Code M MA068; HLDA V: WS Code NK NK51
Immunogen	Human granulocytes

Concentration	1 mg/ml
Preparation	Purified by protein-A affinity chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.