

Technical Data Sheet

Product

Anti-Hu CD25 Biotin

Cat. Number/Size

1B-218-C025

0.025 mg

1B-218-C100

0.1 mg

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Antigen	CD25
Clone	MEM-181
Format	Biotin
Reactivity	Human
Application	FC (QC tested)
Application details	Flow cytometry: Recommended dilution: 1-2 µg/ml.
Isotype	Mouse IgG1
Specificity	The antibody MEM-181 reacts with an extracellular epitope of CD25 (Interleukin-2 receptor alpha chain), a 55 kDa type I transmembrane glycoprotein expressed on activated B and T lymphocytes, activated monocytes/macrophages and on CD4+ T lymphocytes (T regulatory cells); it is lost on resting B and T lymphocytes.
Other names	IL-2R α chain, IL-2-RA, IL2-RA, p55, TAC antigen, IDDM10, TCGFR
Workshop	HLDA VI: WS Code NL N-L024
Immunogen	PHA-activated peripheral blood leucocytes
Entrez Gene ID	3559
Gene name	IL2RA
NCBI Full Gene Name	interleukin 2 receptor subunit alpha
UniProt ID	P01589

Concentration	1 mg/ml
Preparation	Purified antibody is conjugated with biotin LC-NHS ester under optimum conditions and unconjugated antibody and free biotin are removed by size-exclusion chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.